

Sensory quality of custard tart as affected by varying levels of Mabolo (*Dispyros blancoi* A. DC) Flesh

Lorraine P. Baclayon¹, Julious B. Cerna³ and Lynette C. Cimafranca^{2*}

ABSTRACT

Submitted: 1 October 2019 | Accepted: 9 June 2020

Mabolo (*Diospyros blancoi* A. DC) is an underutilized fruit in the Philippines. To add value and maximize the utilization of this nutritional fruit, the potential of mabolo for the production of highly saleable baked products, such as tarts, was explored. Specifically, the study aimed to formulate a custard tart filling with mabolo flesh as a flavorant using a single factor experiment arranged in a Completely Randomized Design (CRD). Six levels of mabolo flesh (0, 15, 30, 45, 60, & 75% w/w) were used. The sensory attributes for color, taste, aroma, texture, flavor, and general acceptability, were evaluated. The optimum formulation of the tart filling was determined using Regression analysis.

Results showed that mabolo custard tart filling had yellowish-brown to brown color, moderately sweet to sweet taste, and soft to slightly fibrous texture. The mabolo fruit aroma and flavor in the tarts ranged from having none to perceptible. The Analysis of Variance revealed that different levels of mabolo flesh cause a significant effect on the color ($p < 0.01$). The optimum formulation and the recommended level of mabolo flesh in custard tart filling production was 80.54% w/w.

Keywords: tart filling, mabolo, sensory evaluation, optimization, tart

INTRODUCTION

Diospyros blancoi A. DC is an indigenous fruit belonging to the Ebenaceae (Ebony) family and is locally called Mabolo (Pobar 2013, Hung et al 2016, Yadav et al 2018). The medium-sized fruit has a skin that turns purple or maroon when ripe and is covered with velvety hairs. The fruit is nutritiously good with a slightly sweet taste and dry texture (Pobar 2013).

¹Department of Food Science and Technology, Visayas State University, Baybay City, Leyte

*Corresponding Author. Address: Department of Food Science and Technology, Visayas State University, Baybay City, Leyte; Email: lynette.cimafranca@vsu.edu.ph
DOI: 10.32945/atr4227.2020

Many of the macro and micronutrients necessary for humans are present in ripe mabolo fruit. Consuming fruits can provide carbohydrates, protein, fiber, and energy. Micronutrients such as vitamin C, vitamin E, sodium, potassium, iron, arsenic (0.01 mg kg^{-1}), calcium, folic acid, pantothenic acid, malic acid, magnesium, phosphorus, zinc, and tannin acid are among the identified compounds contained in mabolo fruits (Hung et al 2016, Haque et al 2009). Mabolo fruit contains various bioactive compounds such as alkaloids, flavonoids, tannins, terpenoids, and essential oils (Maridass et al 2008), which are reported to possess preventive and medicinal properties (Akter et al 2015 as cited by Yadav et al 2018). Despite these beneficial health components in mabolo fruit, utilization of the commodity remains limited and it has minimum to almost no economic use. This may be because the pulp of the mabolo fruit emits a strong cheesy odor when opened possibly deterring consumers. It is often presumed that the fruit has no value; hence no one is interested in the cultivation, enhancement, and propagation of the species. Innovative efforts to transform underutilized and disregarded fruit into a product with high marketability and value are necessary to raise the crop's value. One of the products that are capable of masking the unpleasant and foul cheesy odor is the incorporation of the fruit flesh into highly saleable bakery products (Pobar 2013). One of the easily produced, home baked products visibly seen in the market are tarts.

The tart is defined as a sweet or savory dish with shallow sides and a bottom crust. Usually, tart crusts are made from pastry dough, traditionally made from wheat flour, unsalted butter, cold water, and sugar. Parameters that often affect the quality of tarts are commonly associated with factors such as the quality of the raw materials and the nature of the product. Labor cost, production process, waste of materials, cost of preservation and transportation costs affect the profitability of the product. The production of baked goods requires a number of processing steps between which there exists significant interaction. In terms of quality, the temperature and humidity provide a great impact. The stages of mixing, forming, proofing, baking, and cooling are likewise considered. Aside from the factors mentioned above, the tart filling, which normally dictates variance, may also pose a significant difference in the sensory quality, physical properties and shelf life of the product. Incorporating mabolo flesh, for example, may either cause deteriorative or quality enhancement to the product. Hence, this study aimed to determine the effect of the various levels of mabolo flesh in the filling on the sensory quality of the tart product, as well as to determine the optimum formulation.

MATERIALS AND METHOD

Procurement of Raw Materials

Mabolo fruit was collected from the farms of Abuyog, Leyte. All-purpose flour, sugar, butter, milk, and eggs were purchased at the local market in Baybay City, Leyte.

Preparation of Mabolo Fruit

The *trichomes* (hair) of the mabolo fruit were rubbed off using a brush and washed thoroughly using tap water. The fruits were pared, sliced into 4 to 6 parts,

Sensory quality of custard tart as affected by varying levels of Mabolo

and the seeds were removed. The sliced fruits were steam-blanching for 3 to 5min to stop enzymatic reaction or browning. After steam blanching, the fruit slices were cooled, packed into polyethylene (PE) bags, and stored in the freezer before use.

Preparation of the Mabolo Filling

Frozen mabolo flesh (contained in PE bags) was thawed in running water. It was weighed, and various amounts were added to a custard mixture made of condensed milk (67.07% w/w), eggs (8.67% w/w), butter (7.45% w/w), vanilla (0.87% w/w) and all-purpose flour (15.94% w/w). The different fruit and custard mixtures were cooked in a double boiler. These were then cooled at room temperature for about 5 minutes before being filled into the pastry boat shells (Figure 1).




Figure 1. The process flow chart in the production of mabolo tart

Preparation of the Tart Pastry Shell

Softened butter (43g), eggs (50g), white sugar (250g), and wheat flour (250g) were mixed thoroughly in a clean and sanitized mixing bowl until the tart dough became homogeneous. It was molded into clean, sanitized, and greased tart molders and excess batter was sliced off. Blind-baking at 150°C for 7-8min followed. They were then allowed to cool at room temperature (Figure 1).

Processing of Mabolo Tart

The freshly prepared mabolo filling (20g) was scooped and spread into each of the prepared boat tart shells. They were then placed inside a preheated oven and baked at 180°C for about 20min. The properly baked tarts were removed from the oven and cooled on a rack. After cooling, the tarts were packed in polyethylene bags and stored in an airtight container before analysis.

Experimental Design

A single factor experiment arranged in Completely Randomized Design (CRD) was used in this study. The experiment employed six different levels of mabolo flesh in the production of tart fillings. Table 1 presents the various levels of mabolo flesh that was used in the study.

Table 1. Varying levels of mabolo flesh in tart filling production

| Treatment | Mabolo Flesh (% w/w) |
|-----------|-------------------------|
| 1 | 0 |
| 2 | 15 |
| 3 | 30 |
| 4 | 45 |
| 5 | 60 |
| 6 | 75 |

No. of replication = 2

Sensory Evaluation

The mabolo tart filling was subjected to sensory evaluation to determine the effect of mabolo flesh on the sensory attributes of the product. Color, aroma, taste, flavor, texture and overall acceptability were the sensory attributes evaluated using descriptive scoring and 9-point Hedonic scale. The evaluation was done by 64 sensory panelists at the sensory evaluation room of the Department of Food Science and Technology (DFST), Visayas State University (VSU), Visca, Baybay City, Leyte.

Statistical Analysis

Data were statistically analyzed using a one-way analysis of variance (ANOVA) to establish significant differences between mean values at significance level using

Sensory quality of custard tart as affected by varying levels of Mabolo

Microsoft Excel version 2010. Significantly different results were further analyzed using Tukey's HSD Test to determine further which groups in the samples differ significantly.

The determination of the optimum formulation was done using Regression Analysis in 2nd-degree polynomial using Microsoft Excel™. Point of the lines that intersected with each other was identified as the candidate optimum.

RESULTS AND DISCUSSIONS

Sensory Evaluation

Sensory analysis of food is examined with the human senses. It determines the organoleptic properties of the product and the enjoyment of the consumers towards the products. Sensory testing aims to describe the product's quality characteristics.

Effect on Color

The color of the product often depends on the ingredients used. The color of the mabolo tart filling was brown, as described by 31.5% of the respondents, while 22.7% described it as having a lighter color (light brown). The mean acceptability of tarts without mabolo in the filling (T_1) and those with 15% mabolo flesh (T_2) were 7.67 and 7.50, respectively (Table 3). The lowest color acceptability rating was perceived for tarts with 45% mabolo flesh (T_4) (7.07). However, it should be noted that the forenamed treatment is not significantly different from the rest of the tart products (except T_1) (Table 3). Mean acceptability values were high, which all correspond to 'like moderately' on the 9-point Hedonic scale.

With reference to judges' perception towards individual treatment samples, the panelists described the color of tart without mabolo flesh (T_1) as yellowish-brown, while the color of tart containing 75% mabolo flesh (T_6) was described as brown (Table 2). The flesh of raw mabolo is cream, but sugar in the presence of heat results in chemical reactions such as caramelization and Maillard browning reaction (De Rovira 2008), thereby creating a greater degree of browning at higher levels of mabolo flesh over the control (T_1).

Effect on Aroma

The main aroma compounds in the intact mabolo fruit, pulp, and peel are esters and α -farnesene, according to Hung et al (2016). The perception of the aroma depends on the concentration of these volatiles in the product. In the experiment, the treatment samples with mabolo flesh were described as having a slightly perceptible mabolo fruit aroma (T_2 to T_6). The mean aroma acceptability of the treatment samples for mabolo filling ranges from 7.20- 7.57 (T_3 & T_5), which do not significantly differ from the rest of the tart samples (Table 3). This is equivalent to 'like moderately' in the 9-point Hedonic scale. Increasing mabolo flesh to 30% decreases the aroma acceptability and raises at a 60% level of mabolo flesh (Table 3). Although ANOVA revealed that the utilization of mabolo flesh in the filling causes no significant effect on the aroma acceptability, it is interesting to note that increasing further to 75% could cause acceptability values for the aroma to decline.

Table 2. Quality descriptions of the sensory attributes of tart filling with different levels of mabolo flesh

| TRT | Levels of Mabolo Flesh (%) | Color | Aroma | Flavor | Taste | Texture |
|-----|----------------------------|-----------------|--|--|------------------|------------------|
| 1 | 0 | Yellowish-brown | Absence of mabolo flesh aroma | Absence of mabolo flesh flavor | Moderately sweet | Soft |
| 2 | 15 | Creamy brown | Slightly perceptible mabolo fruit | Slightly perceptible mabolo fruit flavor | Sweet | Soft |
| 3 | 30 | Light brown | Slightly perceptible mabolo fruit aroma | Slightly perceptible mabolo fruit flavor | Sweet | Firm |
| 4 | 45 | Brown | Slightly perceptible to perceptible mabolo fruit aroma | Slightly perceptible mabolo fruit flavor | Sweet | Soft |
| 5 | 60 | Brown | Perceptible mabolo fruit aroma | Perceptible mabolo fruit flavor | Sweet | Soft |
| 6 | 75 | Brown | Slightly perceptible to perceptible mabolo fruit aroma | Perceptible mabolo fruit flavor | Sweet | Slightly fibrous |

9TRT= treatment; n=64

Sensory quality of custard tart as affected by varying levels of Mabolo

Table 3. Response means of the different sensory attributes for mabolo tart filling

| Treatment | Level of Mabolo Flesh (%) | Color* | Aroma ^{ns} | Taste ^{ns} | Flavor ^{ns} | Texture ^{ns} | General Acceptability* |
|-------------|------------------------------|--------------------|---------------------|---------------------|----------------------|-----------------------|------------------------|
| 1 | 0 | 7.67 ^a | 7.47 | 7.63 | 7.47 | 7.63 | 7.73 ^{ab} |
| 2 | 15 | 7.50 ^{ab} | 7.33 | 7.40 | 7.33 | 7.47 | 7.63 ^{ab} |
| 3 | 30 | 7.30 ^{ab} | 7.20 | 7.43 | 7.20 | 7.70 | 7.83 ^{ab} |
| 4 | 45 | 7.07 ^b | 7.43 | 7.47 | 7.43 | 7.30 | 7.90 ^a |
| 5 | 60 | 7.23 ^{ab} | 7.57 | 7.50 | 7.57 | 7.40 | 7.77 ^{ab} |
| 6 | 75 | 7.37 ^{ab} | 7.27 | 7.37 | 7.27 | 7.30 | 7.47 ^b |
| Mean | | 7.36 | 7.38 | 7.47 | 7.38 | 7.47 | 7.72 |

Means with a common letter within a column are not significantly different at 5% level of significance based on Tukey's HSD. 9-point Hedonic Rating Scale: 1=disliked extremely, 2=dislike very much, 3=dislike moderately, 4=dislike slightly, 5=neither like nor dislike, 6=like slightly, 7=like moderately, 8=like very much, 9=like extremely)

Effect on Flavor

According to Figoni (2008), flavor perception depends on many factors related to the product being evaluated as well as to the person doing the evaluation. One of the factors specifically identified by the forenamed author, are the ingredients used in baking. Aside from the complex flavors produced during baking as a result of chemical reactions on carbohydrates, sugars and fats in the crust, the expression of the varying levels of mabolo flesh in the filling on the flavor of the product was assessed.

In terms of flavor, the sensory panelists were able to perceive mabolo flavor in the product except for T₁ (absence of mabolo flesh). Product with higher levels of mabolo flesh had a high perceptibility of mabolo flavor (Table 3). T₄, T₅ and T₆ are distinctively described as having a very perceptible mabolo flavor as shown in Table 2. In other words, the mabolo flavor becomes profound in tarts added with 45, 60, and 75% w/w mabolo flesh. This is due to aromatic compounds, which are present in the mabolo pulp at more concentrated levels, being perceived by retro nasal olfaction during consumption. The condensed milk also contributed to the flavor because of the unique flavor brought about by caramelization and the Maillard reaction during the heating stage of the milk (De Rovira, 2008). The mabolo fruit's characteristic flavor similar to bananas and overripe apple mixed together (Marketman 2004) also contributed to flavor of the product.

Although there was a distinguishing difference in the flavor descriptions of the filling, the ANOVA discloses that varying levels of mabolo flesh do not significantly affect the flavor acceptability of the tart (Table 3). The acceptability values that range from 7.33 to 7.63 are closely comparable, and they all fall at 'like moderately' category of the 9-point Hedonic scale.

Effect on Taste

The mabolo filling samples were described by the panelists as having a sweet taste on treatment samples with 15%, 30%, 45%, 60% and 75% (w/w) mabolo flesh (T₂ to T₆), whereas, 0% level of mabolo flesh was perceived with moderate sweetness (Table 2). The moderately perceived sweetness in T₁ was due purely to the condensed milk, whereas increasing the levels of mabolo flesh resulted in the decreasing sweetness of the product.

The range of acceptability values was 7.37 to 7.63, was within the 'like moderately' category of the 9-point Hedonic scale (Table 3). The ANOVA, however, implied that mabolo flesh at its highest level of 75% had no consequential effect on the acceptability rating for taste.

Effect on Texture

The texture descriptions of the mabolo filling were soft, firm, and slightly fibrous (Table 2). Softness of the filling was perceived by the panelists at treatment samples T₁, T₂, T₄, and T₅, 'firm' for T₃, and 'slightly fibrous' for T₆ (Table 2). Mabolo flesh inherently possesses fibrous materials that become evident at the highest level of mabolo flesh in the product, thereby described as such.

In terms of acceptability for texture, T_1 , T_2 , T_3 showed higher mean acceptability values of 7.63, 7.47, and 7.70, respectively. On the other hand, 7.30, 7.40, and 7.30 are the ratings for T_4 , T_5 , and T_6 , which had slightly lower mean acceptability values. Both groups of values depict an interesting trend at the increasing level of mabolo flesh, but it shall be noted that these values are not significantly different with each other (Table 3). This means that when increasing the level of mabolo flesh up to 75%, texture acceptability was statistically the same as having none. In addition, it was still liked moderately by the panelists.

Effect on General Acceptability

General acceptability is the overall acceptability of the product. T_4 got the highest mean acceptability of 7.90. This acceptability score was statistically revealed not significantly different from T_1 to T_5 , as well as T_6 , but it was statistically different from T_6 (Table 3). Regardless of statistical results, these values correspond to 'like moderately' on the 9-point Hedonic scale. Further, ANOVA conveyed that treatment samples with added mabolo flesh of up to 75% level in the filling were not significantly different from the control.

Optimum Level of Mabolo Flesh in Mabolo Tart Filling

Figure 2 shows that among all of the sensory attributes (from the regression), only the color and general acceptability were correlated with the changes in the levels of mabolo flesh. This is the reason why Figure 3 highlighted the critical levels of mabolo that may be expected as an optimum formulation. At 80.54% (w/w) levels of mabolo flesh, color, and general acceptability intersected. Thus, it was discerned that this level is the optimum level of mabolo flesh in the mabolo tart filling formulation.




Figure 2. Superimposed graph of all sensory attributes for mabolo filling

Sensory quality of custard tart as affected by varying levels of Mabolo




Figure 3. Optimum formulation of mabolo filling

CONCLUSION

Mabolo flesh significantly affects the color of mabolo tart filling, while the rest of the sensory quality parameters (aroma, taste, flavor, texture, & general acceptability) were not significantly affected by the addition of mabolo flesh to the product. Tarts having 75% mabolo flesh in the filling were significantly different from the tarts with 45% mabolo flesh (in terms of general acceptability), while the control was found statistically to have the same general acceptability as samples that had mabolo flesh in the filling. The optimum formulation was 80.54% w/w level of mabolo flesh in the filling.

REFERENCES

- De Rovira D Sr. 2008. Dictionary of flavors (2nd edn). Blackwell Publishing, USA
- Figoni PI. 2010. How Baking Works: Exploring the Fundamentals of Baking Science. John Wiley & Sons, Inc., Hoboken, New Jersey
- Haque MN, Saha BK, Karim MR & Bhuiyan MNH. 2009. Evaluation of Nutritional and Physico-chemical Properties of Several Selected Fruits in Bangladesh. *Bangladesh Journal of Scientific and Industrial Research* 44(3):353-358
- Hung SF, Roan SF, Chang TL, King HB & Chen IZ. 2016. Analysis of aroma compounds and nutrient contents of mabolo (*Diospyros blancoi* A. DC) is an ethnobotanical fruit of Austronesian Taiwan. *Journal of Food and Drug Analysis* 24(1):83-89
- Maridass MG, Ghanthikumar S & Raju G. 2008. Preliminary phytochemical- analysis of *Diospyros Species*. *Ethnobotanical Leaflets* 12:868-72
- Marketman. 2004. MARKETMANILA.COM. <http://www.marketmanila.com/archives/mabolo-kamagong>

- Pobar RA. 2013. Enhancing the use of value-added products from underutilized fruit of the endangered mabolo (*Diospyros blancoi*). *International Journal of Environmental and Rural Development* (4-1):100-105
- Yadav M, Srilekha K, Mrunal BD & Maheswari KU. 2018. Potential health benefit of underutilized fruits: A review. *Journal of Pharmacognosy and Phytochemistry* 7(5):1417-1420